

The majority of men and women who have a chlamydia infection don't know they have it because they have no symptoms and feel completely well.



Chlamydia is easy to test for and treat, so talk to your doctor about a test today.

### Further information

You can find further information about chlamydia at [www.mshc.org.au](http://www.mshc.org.au).

# CHLAMYDIA

## What is Chlamydia?



VCS Pathology acknowledges the support of the Victorian Government  
Copyright Notice © 2018 Victorian Cytology Service Limited [ACN 609 597 408]  
These materials are subject to copyright and are protected by the Copyright Laws of Australia. All rights are reserved.  
Any copying or distribution of these materials without the written permission of the copyright owner is not authorised.

**VCS Pathology**  
PO Box 178, Carlton South, Victoria  
Telephone: [03] 9250 0300 Fax: [03] 9349 1977  
Website: [www.vcs.org.au](http://www.vcs.org.au)

Corp-Mkt-Pub-3 V7



VCS Pathology

## What is Chlamydia?

Chlamydia is a common sexually transmissible infection (STI). If someone has chlamydia they can pass it on to another person during unprotected sex.

Chlamydia can be carried in the:

- cervix (entrance to the uterus at the top of the vagina)
- urethra (the tube which carries urine in men and women)
- rectum (end of large bowel just before anus)

## How do I know if I have Chlamydia?

The majority of men and women who have a chlamydia infection **don't know** they have it because they have **no** symptoms and feel completely well. Only a small number of people with chlamydia have symptoms. These might be:

**In women:**

- abnormal vaginal bleeding (e.g. bleeding when you are **not** having a period, or bleeding after sex)
- vaginal discharge
- pain in the lower abdomen, particularly during sex
- pain or burning when urinating

**In men:**

- discharge from the penis
- stinging, pain or burning when passing urine
- pain in the testicles

## Who should have a Chlamydia test?

- Everyone who has ever had sex and is under the age of 30 should have a chlamydia test every 12 months : it's common in this age group
- Someone with symptoms (see above)
- Men and women with new sexual partners (at least every 12 months)
- Anyone diagnosed with another STI
- Someone who has had sex with a new partner overseas

## How do you test for Chlamydia?


It's simple! If you haven't got any symptoms you just need to pass some urine. You don't need to be examined.

**Your health provider will ask you to go into the toilet and urinate into a jar or onto a small sponge. Not much urine is needed but it is important that it's the first part of the flow.**

**If you have symptoms your health provider will need to examine you. Tests for chlamydia sometimes need to be taken from the vagina or cervix in women.**

When you have your Cervical Screening Test it's a good time to have a chlamydia test as well.

Most tests from men will be on a urine sample.



## Prevention

The best way to prevent infection is to always use condoms with a new partner. And make sure you *both* have a chlamydia test if you want to stop using condoms with your partner.

## How is Chlamydia treated?

Chlamydia is treated with antibiotics---usually just 2 tablets taken together. It is important to treat all sexual partners. Talk to your doctor or nurse about this, and have a look at [www.letthemknow.org.au](http://www.letthemknow.org.au) for some ideas about telling partners. This is very important as they may not know they have a chlamydia infection.

After treatment for a chlamydia infection you should have another test 3 months later.

## What happens if you don't get treatment?

Untreated Chlamydia can cause problems including infertility and pelvic pain.

**In women** untreated chlamydia can cause pelvic inflammatory disease (PID). This is when infection goes into the uterus and the fallopian tubes (the tubes leading from the uterus to the ovaries). This can cause scarring and blockages and could lead to difficulty becoming pregnant.

If the doctor suspects you have PID you will need a longer course of antibiotic treatment---often for a few weeks.

**In men** untreated chlamydia can infect the testicles causing pain, inflammation and sometimes infertility. Again, in this situation a longer course of antibiotics is needed.